

## Sources

## Depression

American Psychiatric Association. Practice guidelines for major depressive disorder in adults. *American Journal of Psychiatry*. 2000; 157(1): S1-S39.

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

Barr L, Goodman WK, Price LH. Physical symptoms associated with paroxetine discontinuation. *Am J Psychiatry*. 1994;151:289.

Beck C. A meta-analysis of predictors of postpartum depression. *Nursing Research*. 1996; 45(5):297-303.

Beck C. Predictors of postpartum depression: an update. *Nursing Research*. 2001; 50(5):275-285.

Bergant A, Heim K, Ulmer H, et al. Early postnatal depressive mood: associations with obstetric and psychosocial factors. *J Psychosom Res*. 1999; 46(4): 391-4.

Borrill J. Detecting and preventing postnatal depression. *Community Nurse*. 1998; 4(3): 19-20.

Breznitz Z, Friedman S. Toddlers' concentration: does maternal depression make a difference? *J Child Psychol Psychiatry*. 1998; 29:267-279.

Brigham and Women's Hospital. *Depression. A guide to diagnosis and treatment*. Boston (MA): Brigham and Women's Hospital; 2001. 9 p.

Chung T, Lau T, Yip A, et al. Antepartum depressive symptomatology is associated with adverse obstetric and neonatal outcomes. *Psychosom Med*. 2001; 63(5): 830-4.

Cox A, Puckering C, Pound A, et al. The impact of maternal depression in young children. *Child Psychol Psychiatry*. 1987; 28:917-928.

Depression Guideline Panel. *Depression in Primary Care: Detection, Diagnosis, and Treatment*. Quick Reference Guide for Clinicians, Number 5. Rockville, MD. U.S. Department of Health and Human Services. Public Health Service, Agency for Health Care Policy and Research. AHCPR Publication No. 93-0552. April 1993.

Epperson C. Postpartum major depression: detection and treatment. *American Family Physician*. 1999; 1-11.

Hobfoll S, Ritter C, Lavin J, et al. Depression prevalence and incidence among inner city pregnant and postpartum women. *J Consult Clin Psychol*. 1995; 63(3): 445-53.

Institute for Clinical Systems Improvement. *Major Depression, Panic Disorder and Generalized Anxiety Disorder in Adults in Primary Care*. Bloomington (MN): Institute For Clinical Systems Improvement (ICSI); 2002.

National Institute for Mental Health. *Depression in Childhood and Adolescents: A fact sheet for physicians*. NIH Publication No. 00-4744. 2000.

- Neter E, Collins N, Lobel M, et al. Psychosocial predictors of postpartum depressed mood in socioeconomically disadvantaged women. *Women's Health*. 1995; 1(1): 51-75.
- Nielsen D, Videbeech P, Hedegaard M, et al. Postpartum depression: identification of women at risk. *BJOG*. 2000; 107(10): 1210-7.
- Pharmacologic treatment of acute major depression and dysthymia. *Ann Intern Med*. 2000; May 2;132(9):738-42.
- Pillai P. Depression: The nurse's role in assessing depression in older people. *Elderly Care*. 1997; 9(3):18-21.
- Seguin L, Potvin L, St-Denis M, et al. Depressive symptoms in the late postpartum among low socioeconomic status women. *Birth*. 1999; 26(3):157-63.
- Spinelli M. Antepartum and postpartum depression. *The Journal of Gender-Specific Medicine*. 1998; 1(2): 33-36.
- Steiner, M. Perinatal mood disorders: position paper. *Psychopharmacol Bull*. 1998; 34(3): 301-6.
- U.S. Preventive Services Task Force. Screening for depression: recommendations and rationale. *Ann Intern Med*. 2002 May 21;136(10):760-4.
- VHA/DOD clinical practice guidelines for the management of major depressive disorder in adults. Washington (DC): Department of Veterans Affairs (U.S.); 2000. Various p.
- Whiffen V, Gotlib I. Infants of postpartum depressed mothers: temperament and cognitive status. *J of Abnormal Psychology*. 1989; 98:274-279.
- Wilson L, Reid A, Midmer D, et al. Antenatal psychosocial risk factors associated with adverse postpartum family outcomes. *CMAJ*, 1996; 154(6):785-799.

### Suicide

- Conwell Y. Suicide in later life: a review and recommendations for prevention. *Suicide and Life Threatening Behavior*. 2001; 31: 32-47.
- Conwell Y. Suicide in elderly patients. In: Schneider L, Reynolds C, Lebowitz B, et al., Depression and treatment in late life: results of NIH consensus development conference. Washington D.C., American Psychiatric Press, 1994; 397-418.
- Gliatto M, Rai A. Evaluation and treatment of patients with suicidal ideation. *American Academy of Family Physicians*. 1999; 1500-1513.
- Holkup P. Evidence based protocol: Elderly suicide. Iowa City (IA): University of Iowa Gerontological Nursing Interventions Research Center, Research Dissemination Core, 2002; Jun. 56p.
- Institute for Clinical Systems Improvement. Major depression, panic disorder and generalized anxiety disorder in adults in primary care. Bloomington (MN): Institute For Clinical Systems Improvement (ICSI); 2002.
- Patterson W, Dohn H, Bird J, Patterson G. Evaluation of suicidal patients: The SAD PERSONS scale. *Psychosomatics*. 1983; 24(4):343-49.

Pillai P. Depression: The nurse's role in assessing depression in older people. *Elderly Care*. 1997; 9(3):18-21.

American Foundation for Suicide Prevention. Available at [www.afsp.org](http://www.afsp.org).

### Substance Abuse

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

CSAT (Center for Substance Abuse Treatment), "A Guide to Substance Abuse Services for Primary Care Physicians," BKD234, U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Treatment Improvement Protocol (TIP Series 24).

Ewing JA. Detecting alcoholism. The CAGE questionnaire. *JAMA*. 1984; 252:1905-7.

Schneider Institute for Health Policy, Brandeis University. Substance Abuse: The Nation's Number One Health Problem. The Robert Wood Johnson Foundation, Princeton, New Jersey; 2001.

### Panic Disorder

American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (4th ed.)*. Washington, D.C.; 1994.

Eaton W, Kramer M, Anthony J, et al. Panic and panic disorder in the United States. *American Journal of Psychiatry*. 1994; 151(3):413-420.

Feldman F. Managing psychiatric disorders in primary care: anxiety. *Hospital Practice*. 2002. Available at [www.hosppractice.com/issues/2000/07/feld.htm](http://www.hosppractice.com/issues/2000/07/feld.htm).

Institute for Clinical Systems Improvement. Major depression, panic disorder and generalized anxiety disorder in adults in primary care. Bloomington (MN): Institute For Clinical Systems Improvement (ICSI); 2002.

Johnson M, et al. Panic disorder in primary care: patients' attributions of illness causes and willingness to accept psychiatric treatment. *Journal of Psychiatric Medicine*. 2000; 30(4) 367-84.

Levinson W, Engel CC. Anxiety. *Behavioral Medicine in Primary Care: A Practical Guide*, Feldman MD, Christensen JF (Eds). Appleton & Lange, Stamford, Conn., 1997, pp 193-211.

Malt E, Berle J, Olafsson S, Lund A, Ursin H. Fibromyalgia is associated with panic disorder and functional dyspepsia with mood disorders. A study of women with random sample population controls. *Journal of Psychosomatic Research*. 2000; 49(5): 285-9.

Miller K. Is collaborative care better in the treatment of panic disorder? American Academy of Family Physicians. 2002. Available at: <http://www.aafp.org/afp/20020401/tips/15.html>. Accessed January 9, 2004.

Narrow W, Rae D, Regier, D. NIMH epidemiology note: prevalence of anxiety disorder. One year estimates calculated from ECA and NCS data. Population estimates based on US Census estimated residential population aged 18 to 54. 1998, unpublished.

Rosenbaum J, Pollack M. *Panic Disorder and Its Treatment*. Marcel Dekker, Inc.; NY: 1998; p.10.

Sheehan D. Treatment of endogenous anxiety with phobic, hysterical, and hypochondriacal symptoms. *Archives of General Psychiatry*. 1980; 37:51-9.

Stein M, Roy-Byrne P, McQuaid J, et al. Development of a brief diagnostic screen for panic disorder in primary care. *Psychosomatic Medicine*. 1999; 61: 359-364.

Yardley L. Relationship between balance system function and agoraphobic avoidance. *Journal of Audiology Medicine*. 1994; 3:48-65.

## PTSD

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

Breslau N, Davis G, Andreski P, et al. Traumatic events and post-traumatic stress disorder in an urban population of young adults. *Archives of General Psychiatry*. 1991; 48(3): 216-22.

Davidson J. Trauma: the impact of post-traumatic stress disorder. *Journal of Psychopharmacology*. 2000; 14(2 Suppl 1): s5-s12.

Feldman F. Managing psychiatric disorders in primary care: Anxiety. *Hospital Practice*. 2002. Available at [www.hosppractice.com/issues/2000/07/feld.htm](http://www.hosppractice.com/issues/2000/07/feld.htm).

Kent JM, Coplan JD, Gorman JM. Clinical utility of the selective serotonin reuptake inhibitors in the spectrum of anxiety. *Biological Psychiatry*. 1998; 44(9): 812.

Lange J, Lange C, Cabaltica R. Primary care treatment of post-traumatic stress disorder. *American Family Physician*. 2000; 62: 1035-46.

Lubin H, Loris M, Burt J, et al. Efficacy of psychoeducational group therapy in reducing symptoms of post-traumatic stress disorder among multiple traumatized women. *American Journal of Psychiatry*. 1998; 155(9): 1172-7.

Marks I, Lovell K, Noshirvani H, et al. Treatment of post-traumatic stress disorder by exposure and/or cognitive restructuring: a controlled study. *Archives of General Psychiatry*. 1998; 55(4): 317-25.

Narrow W, Rae D, Regier, D. NIMH epidemiology note: prevalence of anxiety disorder. One year estimates calculated from ECA and NCS data. Population estimates based on US Census estimated residential population aged 18 to 54. 1998, unpublished.

National Center for Post-traumatic Stress Disorder. Available at [www.ncptsd.com](http://www.ncptsd.com).

Posttraumatic Stress Disorder. American Academy of Child and Adolescent Psychiatry, 1999. Available at [www.aacap.org/publications/factsfam](http://www.aacap.org/publications/factsfam).

## GAD

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

Feldman F. Managing psychiatric disorders in primary care: anxiety. *Hospital Practice*. 2002. Available at [www.hosppract.com/issues/2000/07/feld.htm](http://www.hosppract.com/issues/2000/07/feld.htm).

Institute for Clinical Systems Improvement. Major depression, panic disorder and generalized anxiety disorder in adults in primary care. Bloomington (MN): Institute For Clinical Systems Improvement (ICSI); 2002.

Gliatto M. Generalized anxiety disorder. *American Family Physician*. 2000; 62: 1591-1602.

Levinson W, Engel CC. Anxiety. *Behavioral Medicine in Primary Care: A Practical Guide*. Feldman MD, Christensen JF (Eds). Appleton & Lange, Stamford, Conn., 1997, pp 193-211.

Narrow W, Rae D, Regier, D. NIMH epidemiology note: prevalence of anxiety disorder. One year estimates calculated from ECA and NCS data. Population estimates based on US Census estimated residential population aged 18 to 54. 1998, unpublished.

Rickels K, Schweizer E. The clinical course and long-term management of generalized anxiety disorder. *J Clin Psychopharmacol*. 1990;10:101S-10S.

## OCD

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

Aronson S, Memon M. Obsessive compulsive disorder. *E medicine Journal*. 2002; 3(1).

Feldman F. Managing psychiatric disorders in primary care: anxiety. *Hospital Practice*. 2002. Available at [www.hosppract.com/issues/2000/07/feld.htm](http://www.hosppract.com/issues/2000/07/feld.htm).

Grados M, Scahill L, Riddle MA. Pharmacotherapy in children and adolescents with obsessive-compulsive disorder. *Child Adoles Psych Clinics N Am*. 1999; 8:617-634.

King RA, Scahill L. The assessment and coordination of treatment of children and adolescents with OCD. *Child Adoles Psych Clinics N Am*. 1999; 8:577-597.

Leckman JF. Tourette's syndrome. In: Hollander E, ed. *Obsessive-Compulsive Related Disorders*. Washington, DC: American Psychiatric Press, Inc; 1993.

Leonard HL, Lenane MC, Swedo SE, Rettew DC, Gershon ES, Rapoport JL. Tics and Tourette's disorder: a 2- to 7-year follow-up of 54 obsessive-compulsive children. *Am J Psychiatry*. 1992; 149:1244-1251.

March JS, Leonard HL, Swedo SE. Neuropsychiatry of obsessive-compulsive disorder in children and adolescents. *Compr Ther*. 1995; 21:507-512.

Narrow W, Rae D, Regier, D. NIMH epidemiology note: prevalence of anxiety disorder. One year estimates calculated from ECA and NCS data. Population estimates based on US Census estimated residential population aged 18 to 54. 1998, unpublished.

Steele M, Couturier J. Obsessive compulsive disorder in children and adolescents: assessment in treatment and primary care. *Primary Psychiatry*. 2001; 8(1): 61-66.

## ADHD

American Academy of Pediatrics. Diagnosis and Evaluation of the Child with Attention Deficit/Hyperactivity Disorder (AC0002). *Pediatrics*. 2000; 105(5): 1158-70.

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

Biederman, J. Overview of ADHD. *Contemporary Pediatrics*. 2003; Supplement: 4-5. Institute for Clinical Systems Improvement. Diagnosis and management of ADHD in primary care. Bloomington (MN): Institute for Clinical Systems Improvement, 2000.

Institute for Clinical Systems Improvement. Diagnosis and Management of Attention Deficit Hyperactivity Disorder in Primary Care for School Aged Children and Adolescents: Institute for Clinical Systems Improvement (ICSI); 2003.

Richters J, Arnold L, Jensen P, et al. NIMH collaborative multisite, multimodal treatment study of children with ADHD. *Journal of American Academic Child and Adolescent Psychiatry*. 1995; 34: 987-1000.

Tercyak K, Lerman C, Audrain, J. Association of ADHD symptoms with levels of cigarette smoking in a community sample of adolescents. *J Am Acad Adol Psychiatry*. 2002; 41(7): 799-805.

Wilens T, Faraone S, Biederman J, Gunawardene S. Does stimulant therapy of attention deficit/hyperactivity disorder beget later substance abuse? A meta-analytic review of the literature. *Pediatrics*. 2003; 111(1): 179-85.

## Dementia

American Medical Directors Association (AMDA). Dementia. Columbia (MD): American Medical Directors Association (AMDA); 1998. 32 p.

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (4th ed.)*. Washington, D.C.; 1994.

Cannon B. Dementia vs. Depression. Available at [www.home.epix.net/~tcannon1/depressionvsdementia.htm](http://www.home.epix.net/~tcannon1/depressionvsdementia.htm).

Forester B, Oxman T. Measures to assess the noncognitive symptoms of dementia in the primary care setting. *Journal of Clinical Psychiatry*. 2003; 5(4): 158-163.

Vickery B, Pippenger M. Detection of dementia in primary care. Available at [www.womenshealth.med.ucla.edu/healthcareproviders/dementia.htm](http://www.womenshealth.med.ucla.edu/healthcareproviders/dementia.htm).

## Domestic Violence

AIDS Institute's Clinical Guidelines - "Mental Health Care for People with HIV Infection: HIV Clinical Guidelines for the Primary Care Practitioner." (2001).

American Academy of Pediatrics, Committee on Child Abuse and Neglect. The role of the pediatrician in recognizing and intervening on behalf of abused women. *Pediatrics*. 1998; 101: 1091-1092.

American Medical Association. Diagnostic and Treatment Guidelines on Domestic Violence. Chicago, IL: American Medical Association; 1992.

Borowski I, Ireland M. Parental screening for intimate partner violence by pediatricians and family physicians. *Pediatrics*. 2002; 110(3): 509-516.

Coker A, Smith P, Bethea L, King M, McKeown R. Physical health consequences of physical and psychological intimate partner violence. *Archives of Family Medicine*. 2000; 9(5): 451-7.

Dearwater S, Coben J, Campbell J, Nah S, Glass G, McLoughlin N, Bekemeir, B. Prevalence of intimate partner abuse in women treated at community hospital emergency departments. *Journal of the American Medical Association*. 1998; 280:433-438.

Elliot B, Johnson M. Domestic violence in a primary care setting: patterns and prevalence. *Archives of Family Medicine*. 1995; 4:113-119.

Family Violence Prevention Fund. National Consensus Guidelines on Identifying and Responding to Domestic Violence Victimization in Health Care Settings. 2002.

Mathews, K. Domestic Violence and Primary Care. Department of Pediatrics and Neurology. University of Iowa Hospitals and Clinics. 2003. Available at: [www.vh.org/pediatric/provider/pediatrics/domesticviolence/index.html](http://www.vh.org/pediatric/provider/pediatrics/domesticviolence/index.html). Accessed December 1, 2003.

McCauley J, Kern D, Kolodner K, Dill L, Schroeder A, DeChant H, Ryden J, Bass E, Derogatis L. The “battering syndrome”: prevalence and clinical characteristics of domestic violence in primary care internal medicine practices. *Annals of Internal Medicine*. 1995; 123: 737-746.

National Institute for Justice and Centers for Disease Control. *Prevalence, Incidence and Consequences of Violence Against Women: Findings from the National Violence Against Women Survey*. 1998; Washington, DC: National Institute for Justice and Centers for Disease Control.

New York State Department of Health. Guidelines for integrating domestic violence screening into HIV counseling, testing, referral, and partner notification. Available at: [www.health.state.ny.us/nydoh/rfm/hiv/guide.htm](http://www.health.state.ny.us/nydoh/rfm/hiv/guide.htm). Accessed October 30, 2003.

### Difficult Patients

Christie R, Hoffmaster C. *Ethical Issues in Family Medicine*. Oxford University Press, 1986.

Feltham E. Patients with Personality Disorders. Available at: [www.theberries.ns.ca/Archives/BPDs.html](http://www.theberries.ns.ca/Archives/BPDs.html). Accessed October 30, 2003.

Groves J. Taking care of the hateful patient. *New England Journal of Medicine*. 1978; 298(16):883-887.

Hahn S, Thompson K, Wills T, et al. The difficult doctor-patient relationship: somatization, personality and psychopathology. *J CLin Epidemiology*. 1994; 47(6): 647-57.

Lerner B. Ambulatory Care Syllabus. 1997. Available at: <http://www.cpmc.columbia.edu/whichis/private/aim/34DIFPAT.html>. Accessed January 9, 2004.

- MacLean D. How to manage the hateful patient. *American Medical Directors Association*. 2001; 2(7):1-4.
- Maizes V. Setting limits on demanding patients. *American Family Physician*. 2000; Feb: 1-3.
- Mitchell R. Difficult Patients. 2002. Available at: [www.ttuhscc.edu/pages/fammed/lectures/diffpt/sld003.htm](http://www.ttuhscc.edu/pages/fammed/lectures/diffpt/sld003.htm). Accessed November 20, 2003.
- Noyes R, Holt C, Kathol R. Somatization: Diagnosis and management. *Archives of Family Medicine*. 1995; 4(9): 790-5.
- Schafer S, Nowlis D. Personality disorders among difficult patients. *Archives of Family Medicine*. 1998; 7(2): 126-129.
- Sheehan D. Treatment of endogenous anxiety with phobic, hysterical, and hypochondriacal symptoms. *Archives of General Psychiatry*. 1980; 37:51-9.
- Taub S. Ethics Forum: How do you handle patients' relatives? *American Medical News*. 2000; 3-5.
- White M, Keller V. Difficult clinician-patient relationships. *JCOM*. 1998; 5(5): 32-36.

### Prescribing Antidepressant Medication

- American Psychiatric Association. Practice guidelines for major depressive disorder in adults. *American Journal of Psychiatry*. 2000; 157(1): S1-S39.
- Brigham and Women's Hospital. *Depression. A guide to diagnosis and treatment*. Boston (MA): Brigham and Women's Hospital; 2001. 9 p.
- MacArthur initiative on depression and primary care. 2003. Available at [www.depression-primarycare.org](http://www.depression-primarycare.org).
- Pharmacologic treatment of acute major depression and dysthymia. *Ann Intern Med*. 2000 May 2;132(9):738-42.
- VHA/DOD. Clinical practice guidelines for the management of major depressive disorder in adults. Washington (DC): Department of Veteran Affairs (U.S.); 2000. Various p.